

On Fourth and Higher Moments of Short Exponential Sums Related to Cusp Forms

Anne-Maria Ernvall-Hytönen and Esa V. Vesalainen

Mathematics and Statistics
Åbo Akademi University

Abstract

We obtain upper bounds for the fourth and higher moments of short exponential sums involving Fourier coefficients of holomorphic cusp forms twisted by rational additive twists with small denominators.

1 Introduction

Let us consider a fixed holomorphic cusp form F for $\mathrm{SL}(2, \mathbb{Z})$ of weight $\kappa \in \mathbb{Z}_+$. Then it will have the usual Fourier expansion which we will normalize so that

$$F(z) = \sum_{n=1}^{\infty} a(n) n^{(\kappa-1)/2} e(nz)$$

for all $z \in \mathbb{C}$ with $\Im z > 0$. With this normalization Deligne's estimate [2] says that $a(n) \ll d(n) \ll n^\varepsilon$, for positive integers n , and the Rankin–Selberg estimate [19, 21] says that, for $M \in [1, \infty[$,

$$\sum_{n \leq M} |a(n)|^2 = A M + O(M^{3/5}),$$

where A is a positive real constant only depending on F .

It is of great interest to study exponential sums weighted by Fourier coefficients. Wilton [25] proved essentially square root cancellation for long linear sums and Jutila [16] removed the logarithm in Wilton's estimate leading to the best possible upper bound

$$\sum_{n \leq M} a(n) e(n\alpha) \ll M^{1/2},$$

uniformly true for $M \in [1, \infty[$ and $\alpha \in \mathbb{R}$. The case where α is a reduced fraction h/k with a small denominator k is very interesting, and provides an interesting analogue to the classical problems of studying the error terms in the Dirichlet divisor problem or the circle problem, see e.g. [15]. Jutila [15] proved the pointwise upper bound $\ll k^{2/3} M^{1/3+\varepsilon}$. When $M^{1/10} \ll k \ll M^{5/18}$, this has been improved to $k^{1/4} M^{3/8+\varepsilon}$ in [12, 24], based on short sum estimates from [6].

Jutila [15] also obtained a mean square result analogous to a twisted mean square result for the divisor function in [14], which in turn was in the spirit of earlier work of Cramér [1] for the divisor problem without twists. Crudely speaking, when $k \ll M^{1/2-\varepsilon}$, the size of the sum is proportional to $k^{1/2} M^{1/4}$ on average. More precisely,

$$\int_M^{2M} \left| \sum_{n \leq x} a(n) e\left(n \frac{h}{k}\right) \right|^2 dx = C_F k M^{3/2} + O(k^2 M^{1+\varepsilon}) + O(k^{3/2} M^{5/4+\varepsilon}),$$

where C_F is a positive real constant depending on F only. When $k \ll M^{1/6-\varepsilon}$, it was proved in [24] following [23] and especially [10] that the sum is of the same average order of magnitude in the sense of fourth moments also:

$$\int_M^{2M} \left| \sum_{n \leq x} a(n) e\left(n \frac{h}{k}\right) \right|^4 dx = C'_F k^2 M^2 + O(k^{11/4} M^{15/8+\varepsilon}) + O(k^{13/6} M^{23/12+\varepsilon}),$$

where again C'_F is a positive real constant only depending on the underlying cusp form.

We are interested here in the properties of the short linear sums

$$\sum_{M \leq n \leq M+\Delta} a(n) e(n\alpha),$$

where $M \in [1, \infty[$, $\Delta \in [1, M]$ and $\alpha \in \mathbb{R}$. The best known upper bounds for such sums are due to the first author and Karppinen [6] with a minor improvement by Jääsaari and the second author [12]. We will specifically study the case of a rational α with a small denominator. The study of these short exponential sums is a natural analogue of short interval considerations of error terms in classical analytic number theory. We note that estimates for short sums can also sometimes be used to reduce smoothing error in other arguments, as is done in [16, 3, 6, 24, 12].

Jutila [13] considered the mean square of the error term in the Dirichlet divisor problem in short intervals. The method also applies to short sums of

Fourier coefficients with $\Delta \ll M^{1/2}$, leading to square root cancellation on average, see Ivić [9] and Wu and Zhai [26]. The second moment of short sums of Fourier coefficients with rational additive twists was studied in [4, 24]. The square root cancellation still holds on average as long as $k \ll \Delta^{1/2-\varepsilon}$. More precisely, when $1 \leq \Delta \ll M^{1/2}$ and $k \ll \Delta^{1/2-\varepsilon}$, we have

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^2 dx \ll M \Delta.$$

The second moment of longer short sums was estimated in [5]. The moment estimates give rise to the conjecture that

$$\sum_{M \leq n \leq M+\Delta} a(n) e\left(n \frac{h}{k}\right) \ll \min(\Delta^{1/2} M^\varepsilon, k^{1/2} M^{1/4+\varepsilon}).$$

Ivić [9] considered the fourth moment of the error term in the Dirichlet divisor problem in short intervals and obtained the expected upper bound when the interval was not too short. Wu and Zhai [26] observed that the same technique works for sums of Fourier coefficients. Our first goal here is to consider the fourth moment of short exponential sums with rational additive twists with small denominators in the spirit of [9]. Tanigawa and Zhai [22] extracted a main term in the case of divisor function, but we do not attempt this.

Our second goal is to estimate general higher moments through large value estimates. This follows the consideration of large values of the error term in the Dirichlet divisor problem in short intervals in [11] and the study of higher moments of rationally additive twisted moments of long sums of holomorphic cusp form coefficients in [24], which in turn followed similar study for the moments of the error term in the Dirichlet divisor problem in [8].

2 Notation

We use standard asymptotic notation. If f and g are complex-valued functions defined on some set, say Ω , then we write $f \ll g$ to signify that $|f(x)| \leq C |g(x)|$ for all $x \in \Omega$ for some implicit constant $C \in \mathbb{R}_+$. The notation $O(g)$ denotes a quantity that is $\ll g$, and $f \asymp g$ means that both $f \ll g$ and $g \ll f$. The letter ε denotes a positive real number, whose value can be fixed to be arbitrarily small, and whose value can be different in different instances in a proof. All implicit constants are allowed to depend on ε , on

the implicit constants appearing in the assumptions of theorem statements, and on anything that has been fixed. When necessary, we will use subscripts $\ll_{\alpha, \beta, \dots}$, $O_{\alpha, \beta, \dots}$, etc. to indicate when implicit constants are allowed to depend on objects α, β, \dots .

The numbers $a(1), a(2), \dots$ will always denote the Fourier coefficients of a fixed holomorphic cusp form F of even weight $\kappa \in \mathbb{Z}_+$ for the full modular group $\mathrm{SL}(2, \mathbb{Z})$. The Fourier coefficients are normalized so that the Fourier expansion of the cusp form is

$$F(z) = \sum_{n=1}^{\infty} a(n) n^{(\kappa-1)/2} e(nz)$$

for $z \in \mathbb{C}$ with $\Im z > 0$. All implicit constants are allowed to depend on F .

The function $w(x)$ is a particular kind of smooth weight function, the details of which are given in Definition 11 below. To be precise, all implicit constants are allowed to depend on the L^∞ -norms of w and all its derivatives.

When splitting summation ranges dyadically, we will write

$$\sum_{\substack{L \leq N/2, \\ \text{dyadic}}} \dots,$$

when the summation over L is to be over the values $N/2, N/4, N/8, \dots$, where N is a positive real. These sums will always be finite because the summands will vanish identically for small L . Analogous notation will also be used for various subsums of dyadic sums.

When $h \in \mathbb{Z}$ and $k \in \mathbb{Z}_+$ are coprime, then \bar{h} denotes an integer such that $h\bar{h} \equiv 1 \pmod{k}$.

3 The Results

Let us fix a holomorphic cusp form F of an even weight $\kappa \in \mathbb{Z}_+$ for the full modular group $\mathrm{SL}(2, \mathbb{Z})$. Then F has a Fourier expansion

$$F(z) = \sum_{n=1}^{\infty} a(n) n^{(\kappa-1)/2} e(nz),$$

where, as usual, $z \in \mathbb{C}$ with $\Im z > 0$. Our main theorem on fourth moments is as follows.

Theorem 1. Let $M \in [1, \infty[$, let $\Delta \in [1, M]$, and let $h \in \mathbb{Z}$ and $k \in \mathbb{Z}_+$ be coprime. If $k \ll M^{-1/2} \Delta$ and $k \ll M^{1/4}$, then

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^4 dx \ll k^2 M^{2+\varepsilon}.$$

If $k \gg M^{-1/2} \Delta$ and $k \ll M^{-1/4} \Delta^{2/3}$, then we have

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^4 dx \ll M^{1+\varepsilon} \Delta^2.$$

In the proof we shall mostly, but not entirely, argue analogously to the proof of Theorem 4 in [9].

We wish to detect cancellation in higher moments of short exponential sums. For this purpose, we will estimate the rarity of large values of short exponential sums as follows.

Theorem 2. Let $M, V \in [1, \infty[$, let $\Delta \in [1, M]$, let $\delta \in \mathbb{R}_+$ be fixed, and let h and k be coprime integers with $1 \leq k \leq M$, and assume that $k M^{2\delta} \ll V \ll k M^{1/2+\delta}$. Let $x_1, x_2, \dots, x_R \in [M, 2M]$, where $R \in \mathbb{Z}_+$, and assume that $|x_i - x_j| \geq V$ for $i, j \in \{1, 2, \dots, R\}$ with $i \neq j$. Fix an exponent pair $\langle p, q \rangle \in]0, 1/2] \times [1/2, 1]$. If

$$\sum_{x_i \leq n \leq x_i+\Delta} a(n) e\left(n \frac{h}{k}\right) \gg V$$

for each $i \in \{1, 2, \dots, R\}$, and $k^{2/3} \Delta^{2/3} M^{-1/3+\delta} \ll V \ll \Delta M^\delta$, then

$$R \ll k^2 M^{1+7\delta} \Delta^2 V^{-5} + k^{2q/p} \Delta^{2+2/p} M^{1+q/p+\delta(6+5/p+2q/p)} V^{-2q/p-4-3/p}.$$

Remark. Naturally, the sums in question are always $\ll \Delta M^\varepsilon$ and $\ll \sqrt{M}$, so the condition $V \ll k M^{1/2+\varepsilon}$ will certainly be satisfied in any reasonable application of the result.

Theorem 3. Let $M \in [1, \infty[$ and $\Delta \in [1, M]$, and let h and k be coprime integers with $1 \leq k \leq M$. Let $\alpha, \beta, \gamma \in \mathbb{R}$ be fixed so that

$$\sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \ll k^\alpha \Delta^\beta M^\gamma$$

for $x \in [M, 2M]$. Let $V_0 \in [1, \infty[$ be a parameter such that $k \ll V_0 \ll k^\alpha \Delta^\beta M^\gamma$ and $V_0 \gg k^{2/3} \Delta^{2/3} M^{-1/3}$. Also, let $A \in [2, \infty[$ be fixed and let

$\langle p, q \rangle \in]0, 1/2] \times [1/2, 1]$ be a fixed exponent pair. Then,

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^A dx \ll M^{1+\varepsilon} V_0^A + \Phi + \Psi,$$

where

$$\Phi = \begin{cases} k^2 M^{1+\varepsilon} \Delta^2 V_0^{A-4} & \text{if } A \leq 4, \text{ and} \\ k^{\alpha A - 4\alpha + 2} \Delta^{\beta A - 4\beta + 2} M^{\gamma A - 4\gamma + 1 + \varepsilon} & \text{if } A \geq 4, \end{cases}$$

and

$$\Psi = \begin{cases} k^{2q/p} \Delta^{2+2/p} M^{1+q/p+\varepsilon} V_0^{A-2q/p-3-3/p} & \text{if } A \leq 2q/p + 3 + 3/p, \\ k^{2q/p} \Delta^{2+2/p} M^{1+q/p+\varepsilon} (k^\alpha \Delta^\beta M^\gamma)^{A-2q/p-3-3/p} & \text{otherwise.} \end{cases}$$

It is of course not immediately obvious what exactly this implies. Possible interesting choices for $\langle \alpha, \beta, \gamma \rangle$ are the estimate via absolute values $\langle 0, 1, \varepsilon \rangle$ made possible by Deligne's famous work [2], the estimate for short sums $\langle 0, 1/6, 1/3 + \varepsilon \rangle$ due to [6, Theorem 5.5] which holds when $\Delta \ll M^{2/3}$ [12, Theorem 3], the classical pointwise estimate $\langle 2/3, 0, 1/3 + \varepsilon \rangle$ [15, Corollary on p. 30], as well as the improved pointwise bound $\langle 1/4, 0, 3/8 + \varepsilon \rangle$ which holds for $M^{1/10} \ll k \ll M^{1/4}$ [24, Theorem 1] as well as for $M^{1/4} \ll k \ll M^{5/18}$ [12, Corollary 5].

Let us consider as an example large values of A and k with sums of length $\Delta \asymp M^{5/12}$. We obtain the following upper bound.

Theorem 4. *Let $M \in [1, \infty[$ and let $\Delta \in [1, \infty[$ with $\Delta \asymp M^{5/12}$. Furthermore, let $A \in [11, \infty[$ be fixed, and let h and k be coprime integers with k positive and assume that $M^{1/9} \ll k \ll M^{7/18}$. Then*

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^A dx \ll k^2 M^{11/6+29(A-4)/72+\varepsilon}.$$

As another example, let us consider moments with $A \leq 11$ and $k = 1$ of sums of length $\ll M^{4/9}$. In the following theorem, some of the ranges are treated using similar moment results for long sums from [24].

Theorem 5. *Let $M, \Delta \in [1, \infty[$ with $M^{1/5} \ll \Delta \ll M^{4/9}$ and let $A \in [4, 11]$*

be fixed. Then we have

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) \right|^A dx \ll \begin{cases} M^{A/11+1+\varepsilon} \Delta^{6A/11} & \text{when } A \leq 8 \text{ and } \Delta \ll M^{7/24}, \\ M^{A/4+1+\varepsilon} & \text{when } A \leq 8 \text{ and } \Delta \gg M^{7/24}, \\ M^{A/11+1+\varepsilon} \Delta^{6A/11} & \text{when } A \geq 8 \text{ and } \Delta \ll M^{4/9-11/(9A)}, \\ M^{(A+1)/3+\varepsilon} & \text{when } A \geq 8 \text{ and } \Delta \gg M^{4/9-11/(9A)}. \end{cases}$$

4 Some useful theorems, lemmas and corollaries

4.1 The truncated Voronoi identity for cusp forms

As is to be expected, the proofs use a truncated Voronoi type identity for cusp forms. The following is contained in Theorem 1.1 in [15].

Theorem 6. *Let $x \in [1, \infty[$ and $N \in \mathbb{R}_+$ with $1 \ll N \ll x$, and let h and k be coprime integers such that $1 \leq k \leq x$. Then*

$$\sum_{n \leq x} a(n) e\left(n \frac{h}{k}\right) = \frac{k^{1/2} x^{1/4}}{\pi \sqrt{2}} \sum_{n \leq N} a(n) e\left(-n \frac{\bar{h}}{k}\right) n^{-3/4} \cos\left(4\pi \frac{\sqrt{nx}}{k} - \frac{\pi}{4}\right) + O(k x^{1/2+\varepsilon} N^{-1/2}).$$

Strictly speaking, Theorem 1.1 in [15] assumes that $N \geq 1$ instead of $N \gg 1$. However, if $N \in [c, 1[$, where $c \in]0, 1[$ is fixed, then the identity still holds as stated, for the left-hand side is $\ll x^{1/2}$ by the Wilton–Jutila estimate, and the right-hand side reduces to the O -term $O(k x^{1/2+\varepsilon})$.

4.2 Spacing of square roots

The truncated Voronoi identity leads to exponential sums involving cusp form coefficients with square root phase factors. When expanding a fourth power of such sums we obtain summation over quadruples $\langle a, b, c, d \rangle$. Individual terms will have phase factors involving $\sqrt{a} + \sqrt{b} - \sqrt{c} - \sqrt{d}$, and so we will need a result on the spacing of square roots. The following is contained in Theorem 2 of [20].

Theorem 7. *Let $\omega \in]1, \infty[$ be fixed, let $\delta \in \mathbb{R}_+$ and let $L \geq 2$ be an integer. Then the number of quadruples $\langle a, b, c, d \rangle$ of integers with $a, b, c, d \in]L, 2L]$, and*

$$|a^{1/\omega} + b^{1/\omega} - c^{1/\omega} - d^{1/\omega}| < \delta L^{1/\omega},$$

is

$$\ll \delta L^{4+\varepsilon} + L^{2+\varepsilon}.$$

We shall actually use the special case $\omega = 2$ and $\delta = k M^{\varepsilon-1/2} L^{-1/2}$ for some $M \in [1, \infty[$ and $k \in \mathbb{Z}_+$ with $L \ll M$. It is convenient to observe that in fact real values $L \in [2, \infty[$ are admissible, for if L is not an integer, then we may apply Theorem 7 with $\lfloor L \rfloor$ and $\lceil L \rceil$. The quadruples not covered by these two cases must feature $\lfloor L \rfloor + 1$ and at least one of the two numbers $2 \lceil L \rceil - 1$ and $2 \lfloor L \rfloor$, so that there are at most $\ll L^2$ such quadruples. Finally, when L is smaller, say $L \in [1/2, 2]$, then the number of quadruples is certainly $\ll L^4 \ll 1 \ll L^2$. Thus we have access to the following corollary.

Corollary 8. *Let $M \in [1, \infty[$, $L \in [1/2, \infty[$, $\vartheta \in \mathbb{R}_+$ and $k \in \mathbb{Z}_+$. Then the number of quadruples $\langle a, b, c, d \rangle$ of integers with $a, b, c, d \in]L, 2L]$ and*

$$|\sqrt{a} + \sqrt{b} - \sqrt{c} - \sqrt{d}| < k M^{\vartheta-1/2}$$

is

$$\ll L^{7/2+\varepsilon} k M^{\vartheta-1/2} + L^{2+\varepsilon}.$$

4.3 Plain exponential sums

Our large value estimate depends on estimating certain plain exponential sums. We will do so by employing the machinery of exponent pairs. If $\langle p, q \rangle \in [0, 1/2] \times [1/2, 1]$ is known to be an exponent pair, then

$$\sum_{M \leq n \leq M+\Delta} e(A\sqrt{n}) \ll A^p M^{q-p/2} + A^{-1} M^{1/2},$$

for $M \in [1, \infty[$, $\Delta \in [1, M]$, and $A \in \mathbb{R}_+$. A good reference for the theory of exponent pairs is [7].

We also need the following result which allows us to separate Fourier coefficients from the exponential sums. It is a lemma of Bombieri, and appears as Lemma 1.5 in [18].

Theorem 9. *Let H be a complex Hilbert space with inner product $\langle \cdot | \cdot \rangle$ and norm $\|\cdot\|$. Also, let $\xi, \varphi_1, \varphi_2, \dots, \varphi_R \in H$, where $R \in \mathbb{Z}_+$. Then*

$$\sum_{r=1}^R |\langle \xi | \varphi_r \rangle|^2 \leq \|\xi\|^2 \max_{1 \leq r \leq R} \sum_{s=1}^R |\langle \varphi_r | \varphi_s \rangle|.$$

We shall apply this theorem with $H = \mathbb{C}^N$ for some $N \in \mathbb{Z}_+$ with the usual inner product and norm, which for vectors $z = \langle z_1, \dots, z_N \rangle, w =$

$\langle w_1, \dots, w_N \rangle \in \mathbb{C}^N$ are given by

$$\langle z|w \rangle = \sum_{\ell=1}^N \bar{z}_\ell w_\ell \quad \text{and} \quad \|z\|^2 = \sum_{\ell=1}^N |z_\ell|^2.$$

4.4 Exponential integrals

We will need a lemma for estimating exponential integrals. The following is Lemma 6 in [17].

Lemma 10. *Let $a, b \in \mathbb{R}_+$ and $a < b$, let $g \in C_c^\infty(\mathbb{R}_+)$ with $\text{supp } g \subseteq [a, b]$, and let $G_0, G_1 \in \mathbb{R}_+$ be such that*

$$g^{(\nu)}(x) \ll_\nu G_0 G_1^{-\nu}$$

for all $x \in \mathbb{R}_+$ for each nonnegative integer ν . Also, let f be a holomorphic function defined in $D \subset \mathbb{C}$, which consists of all points in the complex plane with distance smaller than $\rho \in \mathbb{R}_+$ from the interval $[a, b]$ of the real axis. Assume that f is real-valued on $[a, b]$ and let $F_1 \in \mathbb{R}_+$ be such that

$$|f'(z)| \gg F_1$$

for all $z \in D$. Then, for all positive integers P ,

$$\int_a^b g(x) e(f(x)) dx \ll_P G_0 (G_1 F_1)^{-P} \left(1 + \frac{G_1}{\rho}\right)^P (b - a).$$

We remark that, when f is holomorphic in $\{z \in \mathbb{C} | \Re z > 0\}$, we may choose ρ so that $\rho \asymp a$. In particular, in our applications of the lemma, we have $a \asymp b \asymp G_1$ and the factor $(1 + G_1/\rho)^P$ is always $\ll_P 1$.

In the proof of the fourth moment estimate, we will introduce to our integrals a smooth weight function w . For definiteness, we define it here:

Definition 11. In the following, w will denote a function in $C_c^\infty(\mathbb{R}_+)$, depending on $M \in [1, \infty[$, taking values only from the interval $[0, 1]$, and satisfying $\text{supp } w \subseteq [M/2, 5M/2]$, $w \equiv 1$ on $[M, 2M]$, and

$$w^{(\nu)}(x) \ll_\nu M^{-\nu}$$

for all $x \in \mathbb{R}_+$, for every $\nu \in \mathbb{Z}_+ \cup \{0\}$.

The following lemma will be used to estimate several exponential integrals:

Lemma 12. Let $n, k \in \mathbb{Z}_+$, let $\Delta \in \mathbb{R}_+$, assume that $n \ll k^2 M \Delta^{-2}$, and write

$$S(n) = \sin \left(2\pi \frac{\sqrt{n}}{k} \left(\sqrt{x + \Delta} - \sqrt{x} \right) \right).$$

Also, let M and $w(x)$ be as in Definition 11. Then

$$\frac{\partial^\nu}{\partial x^\nu} (w(x) S(a) S(b) S(c) S(d)) \ll_\nu \frac{(abcd)^{1/2} \Delta^4}{k^4 M^{2+\nu}}$$

for all $a, b, c, d \in \mathbb{Z}_+$ with $\max(a, b, c, d) \ll k^2 M \Delta^{-2}$ and $x \in \mathbb{R}_+$, for every $\nu \in \mathbb{Z}_+ \cup \{0\}$.

Proof. Since $w(x)$ vanishes outside the interval $[M/2, 5M/2]$, it is enough to consider the case $x \in [M/2, 5M/2]$. Notice first that we have

$$\begin{aligned} \frac{\partial^\nu}{\partial x^\nu} (w(x) S(a) S(b) S(c) S(d)) &= \sum_{\substack{\alpha_1 + \alpha_2 + \alpha_3 \\ + \alpha_4 + \alpha_5 = \nu}} \frac{\nu!}{\alpha_1! \alpha_2! \alpha_3! \alpha_4! \alpha_5!} \\ &\cdot \left(\frac{\partial^{\alpha_1}}{\partial x^{\alpha_1}} S(a) \right) \left(\frac{\partial^{\alpha_2}}{\partial x^{\alpha_2}} S(b) \right) \left(\frac{\partial^{\alpha_3}}{\partial x^{\alpha_3}} S(c) \right) \left(\frac{\partial^{\alpha_4}}{\partial x^{\alpha_4}} S(d) \right) \left(\frac{\partial^{\alpha_5}}{\partial x^{\alpha_5}} w(x) \right), \end{aligned}$$

where the summation is over quintuples $\langle \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5 \rangle$ of nonnegative integers satisfying $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5 = \nu$. Now

$$S(n) = \sin \left(2\pi \frac{\sqrt{n}}{k} (\sqrt{x + \Delta} - \sqrt{x}) \right) \ll \frac{\sqrt{n}}{k} (\sqrt{x + \Delta} - \sqrt{x}) \ll \frac{\sqrt{n} \Delta}{k \sqrt{x}},$$

and when $\alpha \in \mathbb{Z}_+ \cup \{0\}$, we have

$$\frac{\partial^\alpha}{\partial x^\alpha} S(n) \ll_\alpha \frac{\sqrt{n} \Delta}{k x^{(2\alpha+1)/2}}.$$

Putting everything together, we obtain

$$\begin{aligned} \frac{\partial^\nu}{\partial x^\nu} (w(x) S(a) S(b) S(c) S(d)) &= \sum_{\substack{\alpha_1 + \alpha_2 + \alpha_3 \\ + \alpha_4 + \alpha_5 = \nu}} \frac{\nu!}{\alpha_1! \alpha_2! \alpha_3! \alpha_4! \alpha_5!} \\ &\cdot \left(\frac{\partial^{\alpha_1}}{\partial x^{\alpha_1}} S(a) \right) \left(\frac{\partial^{\alpha_2}}{\partial x^{\alpha_2}} S(b) \right) \left(\frac{\partial^{\alpha_3}}{\partial x^{\alpha_3}} S(c) \right) \left(\frac{\partial^{\alpha_4}}{\partial x^{\alpha_4}} S(d) \right) \left(\frac{\partial^{\alpha_5}}{\partial x^{\alpha_5}} w(x) \right) \\ &\ll_\nu \sum_{\substack{\alpha_1 + \alpha_2 + \alpha_3 \\ + \alpha_4 + \alpha_5 = \nu}} \frac{\sqrt{a} \Delta}{k x^{(2\alpha_1+1)/2}} \cdot \frac{\sqrt{b} \Delta}{k x^{(2\alpha_2+1)/2}} \cdot \frac{\sqrt{c} \Delta}{k x^{(2\alpha_3+1)/2}} \cdot \frac{\sqrt{d} \Delta}{k x^{(2\alpha_4+1)/2}} \cdot x^{-\alpha_5} \\ &\ll_\nu \frac{(abcd)^{1/2} \Delta^4}{k^4 M^{2+\nu}}. \end{aligned}$$

□

Before embarking on the proofs of Theorems 1, 2 and 3, we introduce one final lemma on the mean square of the kind of exponential sums which arise from the truncated Voronoi identity.

Lemma 13. *Let $M \in [1, \infty[$, $L \in [1/2, \infty[$ and $T \in [0, \infty[$ with $T \ll M$ and let $h \in \mathbb{Z}$ and $k \in \mathbb{Z}_+$ be coprime. Furthermore, let $w(x)$ be a smooth weight function as in Definition 11. Then we have*

$$\int_{M/2}^{5M/2} w(x) \left| \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) e\left(\pm \frac{2\sqrt{n(x+T)}}{k}\right) \right|^2 dx \ll M L^{-1/2} + L^\varepsilon k M^{1/2+\varepsilon},$$

and if we further assume that $L \ll M^{1-\vartheta} k^{-2}$ for some fixed positive real number ϑ that can be chosen to be arbitrarily small, then we have

$$\int_{M/2}^{5M/2} w(x) \left| \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) e\left(\pm \frac{2\sqrt{n(x+T)}}{k}\right) \right|^2 dx \ll M L^{-1/2}.$$

Proof. Here we expand the square as $|\Sigma|^2 = \Sigma \bar{\Sigma}$ and separate the diagonal terms from the off-diagonal terms, leading to

$$\begin{aligned} & \int_{M/2}^{5M/2} w(x) \left| \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) e\left(\pm \frac{2\sqrt{n(x+T)}}{k}\right) \right|^2 dx \\ & \ll \sum_{L < n \leq 2L} \frac{|a(n)|^2}{n^{3/2}} \int_{M/2}^{5M/2} w(x) dx \\ & \quad + \sum_{L < m < n \leq 2L} \frac{|a(m) a(n)|}{(mn)^{3/4}} \left| \int_{M/2}^{5M/2} w(x) e\left(\pm \frac{2(\sqrt{m} - \sqrt{n})\sqrt{x+T}}{k}\right) dx \right|. \end{aligned}$$

The diagonal terms contribute $\ll M L^{-1/2}$, and by Lemma 10, the off-diagonal terms contribute, for arbitrary $P \in \mathbb{Z}_+$,

$$\ll_P \sum_{L < m < n \leq 2L} \frac{|a(m) a(n)|}{(mn)^{3/4}} \cdot \left(\frac{k M^{-1/2}}{\sqrt{n} - \sqrt{m}} \right)^P \cdot M.$$

When $k M^{-1/2} |\sqrt{n} - \sqrt{m}|^{-1} \ll M^{-\varepsilon'}$, for some constant $\varepsilon' \in \mathbb{R}_+$, the bound above can be made as small as desired by choosing P to be sufficiently large (depending on ε'). Let us now choose $\varepsilon' \in]0, \vartheta/2[$. The condition

$$\frac{k}{M^{1/2} |\sqrt{n} - \sqrt{m}|} \ll M^{-\varepsilon'}$$

holds when $|n - m| \gg k M^{-1/2+\varepsilon'} \sqrt{L}$. Therefore, when $L \ll M^{1-\vartheta} k^{-2}$, we have

$$k M^{-1/2+\varepsilon'} \sqrt{L} \ll k^{1-1} M^{-1/2+\varepsilon'+1/2-\vartheta/2} = o(1),$$

and hence, when $m \neq n$ and so $|m - n| \geq 1$, we have

$$\frac{k}{M^{1/2} |\sqrt{n} - \sqrt{m}|} \ll M^{-\varepsilon'},$$

and thus, in particular, only the contribution from diagonal terms counts when $L \ll M^{1-\vartheta} k^{-2}$.

Let us now estimate the contribution coming from the off-diagonal terms for which $|n - m| \ll k M^{-1/2+\varepsilon'} \sqrt{L}$, when $L \gg M^{1-\vartheta} k^{-2}$. For each value of n , there are $\ll \sqrt{L} k M^{-1/2+\varepsilon'}$ values of m , and for all of these values, we estimate the integral by absolute values. We thus obtain from the off-diagonal terms

$$\ll \sum_{L < n \leq 2L} L^{\varepsilon-3/2} \sqrt{L} k M^{-1/2+\varepsilon} M \ll L^\varepsilon k M^{1/2+\varepsilon'}.$$

□

4.5 Moments of long linear sums

The following is Theorem 2.3 from [24].

Theorem 14. *Let $M \in [1, \infty[$, let us fix an exponent pair $\langle p, q \rangle \in]0, 1/2] \times [1/2, 1]$ satisfying $q \geq (p + 1)/2$, and let h and k be coprime integers with $1 \leq k \ll M^{1/2-\varepsilon}$. Furthermore, let $\alpha, \beta, \gamma, \delta, A \in [0, \infty[$ be fixed exponents so that*

$$\sum_{n \leq x} a(n) e\left(n \frac{h}{k}\right) \ll k^\alpha x^{\beta+\varepsilon}$$

for $x \in [1, \infty[$ and for k satisfying $x^\gamma \ll k \ll x^\delta$. Then, for $M^\gamma \ll k \ll M^\delta$,

$$\int_M^{2M} \left| \sum_{n \leq x} a(n) e\left(n \frac{h}{k}\right) \right|^A dx \ll k^{A/2} M^{A/4+1} + \Phi + \Psi,$$

where

$$\Phi = \begin{cases} k^{\alpha A + 2(1-\alpha)} M^{\beta A + (1-2\beta) + \varepsilon} & \text{if } A \geq 2, \\ k^{A/2+1} M^{A/4+1/2+\varepsilon} & \text{if } A \leq 2, \end{cases}$$

and

$$\Psi = \begin{cases} k^{\alpha A - \alpha - \alpha/p + (1-\alpha)2q/p} M^{\beta A + 1 - \beta - \beta/p + (1-2\beta)q/p + \varepsilon} & \text{if } A \geq 1 + (1 + 2q)/p, \\ k^{A/2 - 1/2 - 1/(2p) + q/p} M^{A/4 + 3/4 - 1/(4p) + q/(2p) + \varepsilon} & \text{if } A \leq 1 + (1 + 2q)/p. \end{cases}$$

5 Proof of Theorem 1

We let $\varepsilon_0 \in \mathbb{R}_+$ be arbitrary. Our goal is to prove an estimate $\ll M^{2+\varepsilon_0} k^2$ or $\ll M^{1+\varepsilon_0} \Delta^2$. Some exponents in the proof will depend on the desired final value of ε_0 . We assume throughout the proof that $k \ll M^{-1/2} \Delta$ and $k \ll M^{1/4}$, or that $k \gg M^{-1/2} \Delta$ and $k \ll M^{-1/4} \Delta^{2/3}$.

We begin by applying the truncated Voronoi identity for cusp form coefficients to get, for $N \in \mathbb{R}_+$ satisfying $1 \ll N \ll M$,

$$\begin{aligned} & \int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^4 dx \\ & \ll \int_M^{2M} \left| k^{1/2} \sum_{n \leq N} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) \left((x+\Delta)^{1/4} \cos\left(4\pi \frac{\sqrt{n(x+\Delta)}}{k} - \frac{\pi}{4}\right) \right. \right. \\ & \quad \left. \left. - x^{1/4} \cos\left(4\pi \frac{\sqrt{nx}}{k} - \frac{\pi}{4}\right) \right) \right|^4 dx + M^{3+\varepsilon} N^{-2} k^4, \end{aligned}$$

where we applied the elementary inequality $|A+B|^4 \ll |A|^4 + |B|^4$, which holds uniformly for all $A, B \in \mathbb{C}$.

Let us now choose N in the following way:

$$N = \begin{cases} M^{1/2} k, & \text{when } k \ll \Delta M^{-1/2}, \\ k^2 M \Delta^{-1}, & \text{otherwise.} \end{cases}$$

Thus, when $k \ll \Delta M^{-1/2}$, we have trivially $N \geq 1$, and we have $N \ll M$ since $k \ll M^{1/2}$. When $k \gg \Delta M^{-1/2}$, we have again trivially $N \gg 1$, and we have $N \ll M$ since $k \ll M^{-1/4} \Delta^{2/3} \ll \Delta^{2/3-1/4} = \Delta^{5/12} \ll \Delta^{1/2}$.

Hence the error from the error term of the truncated Voronoi identity becomes

$$M^{3+\varepsilon} N^{-2} k^4 \ll \begin{cases} M^{2+\varepsilon} k^2, & \text{when } k \ll \Delta M^{-1/2}, \\ M^{1+\varepsilon} \Delta^2, & \text{otherwise.} \end{cases}$$

Since the integrand is nonnegative, we may introduce the weight function $w(x)$ of Definition 11 to the integral involving the main terms from the truncated Voronoi identity, and extend the region of integration to be over the interval $[M/2, 5M/2]$:

$$\int_M^{2M} |\dots|^4 dx \ll \int_{M/2}^{5M/2} w(x) |\dots|^4 dx.$$

Next, we split the sum \sum_n dyadically into $\sum_{L \leq N/2} \sum_{L < n \leq 2L}$, where L ranges over the values $N/2, N/4, N/8, \dots$. There will be $\ll 1 + \log M$ such values of interest, and so we may continue the estimations by applying Hölder's inequality to get

$$\begin{aligned} &\ll (1 + \log M)^3 \sum_{\substack{L \leq N/2 \\ \text{dyadic}}} \int_{M/2}^{5M/2} w(x) \\ &\cdot \left| k^{1/2} \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) \left((x + \Delta)^{1/4} \cos(\dots) - x^{1/4} \cos(\dots) \right) \right|^4 dx, \end{aligned}$$

where of course $(1 + \log M)^3 \ll M^\varepsilon$. The sum over L is split into three parts: those terms with L large, the terms with L so small, that there is very little oscillation, but there is cancellation in the main terms of the truncated Voronoi identity, and the remaining terms in the middle:

$$\sum_{\substack{L \leq N/2 \\ \text{dyadic}}} = \sum_{\substack{L \ll Y \\ \text{dyadic}}} + \sum_{\substack{Y \ll L \ll X \\ \text{dyadic}}} + \sum_{\substack{X \ll L \leq N/2 \\ \text{dyadic}}}.$$

When $k \ll \Delta M^{-1/2}$ we choose

$$Y = 1 \quad \text{and} \quad X = \sqrt{M},$$

and certainly $1 \ll Y \ll X \ll N$. In particular, the first sum over $L \ll Y$ will be empty.

When $k \gg \Delta M^{-1/2}$, we choose

$$Y = k^2 M \Delta^{-2} \quad \text{and} \quad X = \min(\Delta^4 M^{-1} k^{-6}, N).$$

We will trivially have $1 \ll Y \ll N$. Also, we always have $Y \ll \Delta^4 M^{-1} k^{-6}$ since this is equivalent with $k \ll M^{-1/4} \Delta^{3/4}$ and this holds since we have $k \ll M^{-1/4} \Delta^{2/3} \ll M^{-1/4} \Delta^{3/4}$. When $\Delta \gg M^{2/5} k^{8/5}$, we have $X \asymp N$ and the sum over $L \gg X$ will be empty. The rest of the proof consists of working through each of these cases separately.

The high-frequency terms $L \gg X$. For these values of L , we may estimate by the truncated Voronoi identity that for any $x \in [M/2, 5M/2]$ and each $T \in \{0, \Delta\}$,

$$\begin{aligned}
& k^{1/2} (x+T)^{1/4} \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) \cos\left(4\pi \frac{\sqrt{n}}{k} \sqrt{x+T} - \frac{\pi}{4}\right) \\
&= k^{1/2} (x+T)^{1/4} \sum_{n \leq 2L} \dots - k^{1/2} (x+T)^{1/4} \sum_{n \leq L} \dots \\
&= \pi \sqrt{2} \sum_{n \leq x+T} a(n) e\left(n \frac{h}{k}\right) + O(k M^{1/2+\varepsilon} L^{-1/2}) \\
&\quad - \pi \sqrt{2} \sum_{n \leq x+T} a(n) e\left(n \frac{h}{k}\right) - O(k M^{1/2+\varepsilon} L^{-1/2}) \ll k M^{1/2+\varepsilon} L^{-1/2},
\end{aligned}$$

so that

$$\left| k^{1/2} \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) \left((x+\Delta)^{1/4} \cos(\dots) - x^{1/4} \cos(\dots) \right) \right|^2 \ll k^2 M^{1+\varepsilon} L^{-1}.$$

The contribution from the high-frequency terms involving $\sqrt{x+T}$, where $T \in \{0, \Delta\}$, can be estimated by

$$\begin{aligned}
& \ll k^3 M^{3/2+\varepsilon} \sum_{\pm} \sum_{\substack{X \ll L \leq N/2 \\ \text{dyadic}}} L^{-1} \\
& \cdot \int_{M/2}^{5M/2} w(x) \left| \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) e\left(\pm 2 \frac{\sqrt{n}}{k} \sqrt{x+T}\right) \right|^2 dx.
\end{aligned}$$

We may use Lemma 13 to bound the expression on the second line. The contribution coming from this is $\ll M L^{-1/2} + L^\varepsilon k M^{1/2+\varepsilon}$, and if $L \ll M^{1-\varepsilon_0/2} k^{-2}$, then the contribution is $\ll M L^{-1/2}$.

The contribution coming from the diagonal terms $M L^{-1/2}$ is

$$\ll k^3 M^{3/2+\varepsilon} \sum_{\substack{X \ll L \leq N/2 \\ \text{dyadic}}} L^{-1} M L^{-1/2} \ll k^3 M^{5/2+\varepsilon} X^{-3/2}.$$

When $k \ll \Delta M^{-1/2}$, we have $X = M^{1/2}$, and hence the contribution will be

$$\ll k^3 M^{5/2+\varepsilon} M^{-3/4} = k^3 M^{7/4+\varepsilon} \ll k^2 M^{2+\varepsilon},$$

since $k \ll M^{1/4}$. When $k \gg \Delta M^{-1/2}$, we have $X \asymp N$, in which case there are no high-frequency terms to consider, or $X = \Delta^4 M^{-1} k^{-6}$. In the latter case we obtain

$$\begin{aligned} &\ll k^3 M^{5/2+\varepsilon} X^{-3/2} \ll k^3 M^{5/2+\varepsilon} (\Delta^4 M^{-1} k^{-6})^{-3/2} \\ &\ll k^{12} M^{4+\varepsilon} \Delta^{-6} \ll M^{1+\varepsilon} \Delta^2, \end{aligned}$$

since $k \ll \Delta^{2/3} M^{-1/4}$.

Finally, let us compute the contribution of the term $L^\varepsilon k M^{1/2+\varepsilon}$. This term exists only for $L \gg M^{1-\varepsilon_0/2} k^{-2}$, and we thus obtain

$$\begin{aligned} &\ll k^3 M^{3/2+\varepsilon} \sum_{\substack{M^{1-\varepsilon_0/2} k^{-2} \ll L \leq N/2 \\ \text{dyadic}}} L^{\varepsilon-1} k M^{1/2+\varepsilon} \\ &\ll k^4 M^{2+\varepsilon} (M^{1-\varepsilon_0/2} k^{-2})^{\varepsilon-1} \ll M^{1+\varepsilon_0/2+\varepsilon} k^6. \end{aligned}$$

In the case $k \ll \Delta M^{-1/2}$ this is $\ll k^2 M^{2+\varepsilon_0}$ since $k \ll M^{1/4}$. In the case $k \gg \Delta M^{-1/2}$ this is $\ll M^{1+\varepsilon_0} \Delta^2$, provided that $k \ll \Delta^{1/3}$. But this holds since $M^{-1/2} \Delta \ll k \ll M^{-1/4} \Delta^{2/3}$, so that $\Delta \ll M^{3/4}$, and therefore $k \ll M^{-1/4} \Delta^{2/3} \ll \Delta^{-1/3} \Delta^{2/3} \ll \Delta^{1/3}$.

The low-frequency terms $L \ll Y$. Let us recall first that these terms need to be considered only in the case $k \gg \Delta M^{-1/2}$ in which $Y = k^2 M \Delta^{-2}$.

For low-frequency terms we want to get the sums to partially cancel each other, and therefore, we want to replace the factor $(x + \Delta)^{1/4}$ by $x^{1/4}$:

$$\begin{aligned} &k^{1/2} \sum_{L < n \leq 2L} \frac{a(n)}{n^{3/4}} e\left(-n \frac{\bar{h}}{k}\right) ((x + \Delta)^{1/4} - x^{1/4}) \cos\left(4\pi \frac{\sqrt{n(x + \Delta)}}{k} - \frac{\pi}{4}\right) \\ &\ll k^{1/2} L^{1/4} \Delta M^{-3/4} \ll k^{1/2} (k^2 M \Delta^{-2})^{1/4} \Delta M^{-3/4} \ll k M^{-1/2} \Delta^{1/2}. \end{aligned}$$

Hence the total contribution coming from replacing $(x + \Delta)^{1/4}$ by $x^{1/4}$ is

$$\ll M^{1+\varepsilon} (k M^{-1/2} \Delta^{1/2})^4 \ll k^4 M^{-1+\varepsilon} \Delta^2 \ll \Delta^2 M^{1+\varepsilon},$$

which holds since $k \ll M^{-1/4} \Delta^{2/3} \ll M^{2/3-1/4} = M^{5/12} \ll M^{1/2}$. We may now use the elementary trigonometric identity

$$\cos\left(2\xi - \frac{\pi}{4}\right) - \cos\left(2\eta - \frac{\pi}{4}\right) = 2 \sin(\xi - \eta) \cos\left(\xi + \eta + \frac{\pi}{4}\right),$$

which holds for any $\xi, \eta \in \mathbb{R}$. Applying this with

$$\xi = 2\pi \frac{\sqrt{n}}{k} \sqrt{x + \Delta} \quad \text{and} \quad \eta = 2\pi \frac{\sqrt{n}}{k} \sqrt{x},$$

the contribution from the terms with $L \ll Y$ is

$$\begin{aligned}
&\ll k^2 M^{1+\varepsilon} \sum_{\pm} \sum_{\substack{L \ll Y \\ \text{dyadic}}} \\
&\cdot \int_{M/2}^{5M/2} w(x) \left| \sum_{L < n \leq 2L} \frac{a(n)}{n^{3/4}} e\left(-n \frac{\bar{h}}{k}\right) S(n) e\left(\pm \frac{\sqrt{n}}{k} (\sqrt{x+\Delta} + \sqrt{x})\right) \right|^4 dx \\
&\ll k^2 M^{1+\varepsilon} \sum_{\substack{L \ll Y \\ \text{dyadic}}} \sum_{L < a \leq 2L} \sum_{L < b \leq 2L} \sum_{L < c \leq 2L} \sum_{L < d \leq 2L} \frac{|a(a) a(b) \overline{a(c)} a(d)|}{(abcd)^{3/4}} \\
&\cdot \left| \int_{M/2}^{5M/2} w(x) S(a) S(b) S(c) S(d) e\left(\frac{\alpha}{k} (\sqrt{x+\Delta} + \sqrt{x})\right) dx \right|.
\end{aligned}$$

where the factors $S(n)$ are given by

$$S(n) = \sin\left(2\pi \frac{\sqrt{n}}{k} (\sqrt{x+\Delta} - \sqrt{x})\right),$$

the coefficient α is the square root expression

$$\alpha = \sqrt{a} + \sqrt{b} - \sqrt{c} - \sqrt{d},$$

and w is as in Definition 11.

Let us first consider the terms of $\sum_a \sum_b \sum_c \sum_d$ with $\alpha \gg M^{\varepsilon_0/2-1/2} k$. Using Lemma 12, we have, for each $\nu \in \mathbb{Z}_+ \cup \{0\}$,

$$\frac{d^\nu}{dx^\nu} (w(x) S(a) S(b) S(c) S(d)) \ll_\nu \frac{(abcd)^{1/2} \Delta^4}{k^4 M^{2+\nu}}.$$

Therefore, in the terms under consideration, the integral $\int_{M/2}^{5M/2} \dots dx$ may be estimated using Lemma 10 to be, for any $P \in \mathbb{Z}_+$,

$$\begin{aligned}
&\ll_P \frac{(abcd)^{1/2} \Delta^4}{k^4 M^2} (M \alpha M^{-1/2} k^{-1})^{-P} M \ll \frac{(abcd)^{1/2} \Delta^4}{k^4 M^2} M^{1-P\varepsilon_0/2} \\
&\ll (abcd)^{1/2} M^{1-P\varepsilon_0/2}.
\end{aligned}$$

Fixing P to be sufficiently large (depending on ε_0), the contribution from the

terms under consideration will be

$$\begin{aligned}
&\ll k^2 M^{1+\varepsilon} \sum_{\substack{L \ll Y \\ \text{dyadic}}} \sum_a \sum_b \sum_c \sum_d \frac{|a(a) a(b) a(c) a(d)|}{(abcd)^{1/4}} M^{1-P\varepsilon_0/2} \\
&\ll k^2 M^{2+\varepsilon-P\varepsilon_0/2} \sum_{\substack{L \ll Y \\ \text{dyadic}}} L^3 \ll M^{6+\varepsilon-P\varepsilon_0/2} \ll 1.
\end{aligned}$$

Finally, by Corollary 8, the number of terms in the sum $\sum_a \sum_b \sum_c \sum_d$ with $\alpha \ll k M^{\varepsilon_0/2-1/2}$ is

$$\ll L^{7/2+\varepsilon} k M^{\varepsilon_0/2-1/2} + L^{2+\varepsilon},$$

and so we conclude, estimating everything by absolute values, and sine factors by $\sin x \ll x$, that the rest of the low-frequency terms with $L \ll Y$ contribute

$$\begin{aligned}
&\ll k^2 M^{1+\varepsilon} \sum_{\substack{L \ll Y \\ \text{dyadic}}} (L^{7/2+\varepsilon} k M^{\varepsilon_0/2-1/2} + L^{2+\varepsilon}) L^{\varepsilon-3} M (L^{1/2} \Delta M^{-1/2} k^{-1})^4 \\
&\ll k^2 \Delta^4 M^\varepsilon \sum_{\substack{L \ll Y \\ \text{dyadic}}} (L^{5/2+\varepsilon} k^{-3} M^{\varepsilon_0/2-1/2} + L^{1+\varepsilon} k^{-4}) \\
&\ll \Delta^4 M^\varepsilon \left((k^2 M \Delta^{-2})^{5/2+\varepsilon} k^{-1} M^{\varepsilon_0/2-1/2} + (k^2 M \Delta^{-2})^{1+\varepsilon} k^{-2} \right) \\
&\ll \Delta^{-1} M^{2+\varepsilon_0/2+\varepsilon} k^4 + \Delta^2 M^{1+\varepsilon} \ll M^{1+\varepsilon_0} \Delta^2,
\end{aligned}$$

since $k \ll \Delta^{2/3} M^{-1/4} \ll \Delta^{3/4} M^{-1/4}$.

The terms in the middle with $Y \ll L \ll X$. The contribution from the terms with $Y \ll L \ll X$ and involving $\sqrt{x+T}$, where $T \in \{0, \Delta\}$, is

$$\begin{aligned}
&\ll k^2 M^{1+\varepsilon} \sum_{\pm} \\
&\quad \cdot \sum_{\substack{Y \ll L \ll X \\ \text{dyadic}}} \int_{M/2}^{5M/2} w(x) \left| \sum_{L < n \leq 2L} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k} \pm 2 \frac{\sqrt{n}}{k} \sqrt{x+T}\right) \right|^4 dx \\
&\ll k^2 M^{1+\varepsilon} \sum_{\substack{Y \ll L \ll X \\ \text{dyadic}}} \sum_{L < a \leq 2L} \sum_{L < b \leq 2L} \sum_{L < c \leq 2L} \sum_{L < d \leq 2L} \frac{|a(a) a(b) \overline{a(c)} a(d)|}{(abcd)^{3/4}} \\
&\quad \cdot \left| \int_{M/2}^{5M/2} w(x) e\left(2\alpha \frac{\sqrt{x+T}}{k}\right) dx \right|,
\end{aligned}$$

where the coefficient α is again the square root expression

$$\alpha = \sqrt{a} + \sqrt{b} - \sqrt{c} - \sqrt{d}.$$

In those terms of $\sum_a \sum_b \sum_c \sum_d$ in which $\alpha \gg k M^{\varepsilon_0/2-1/2}$, we may estimate the integral $\int_{M/2}^{5M/2}$ by Lemma 10 for any $P \in \mathbb{Z}_+$ by

$$\ll_P (M \alpha k^{-1} M^{-1/2})^{-P} M \ll M^{1-P\varepsilon_0/2}.$$

Thus, these terms contribute, taking P fixed and sufficiently large (depending on ε_0),

$$\ll k^2 M^{1+\varepsilon} \sum_{\substack{Y \ll L \ll X \\ \text{dyadic}}} L^{1+\varepsilon} M^{1-P\varepsilon_0/2} \ll k^2 M^{2+\varepsilon-P\varepsilon_0/2} X^{1+\varepsilon} \ll 1.$$

Finally, the number of terms in $\sum_a \sum_b \sum_c \sum_d$ in which $\alpha \ll k M^{\varepsilon_0/2-1/2}$ is by Corollary 8

$$\ll L^{7/2+\varepsilon} k M^{\varepsilon_0/2-1/2} + L^{2+\varepsilon},$$

and so the contribution from these terms, estimating by absolute values, is

$$\begin{aligned} &\ll k^2 M^{1+\varepsilon} \sum_{\substack{Y \ll L \ll X \\ \text{dyadic}}} (L^{7/2+\varepsilon} k M^{\varepsilon_0/2-1/2} + L^{2+\varepsilon}) L^{\varepsilon-3} M \\ &\ll k^2 M^{2+\varepsilon} \sum_{\substack{Y \ll L \ll X \\ \text{dyadic}}} (L^{1/2} k M^{\varepsilon_0/2-1/2} + L^{-1}). \end{aligned}$$

The contribution from the second term L^{-1} is

$$\ll \begin{cases} k^2 M^{2+\varepsilon} & \text{if } k \ll \Delta M^{-1/2} \\ \Delta^2 M^{1+\varepsilon} & \text{otherwise.} \end{cases}$$

Let us now move to considering the first term. In the case $k \ll \Delta M^{-1/2}$, we have $X = M^{1/2}$, and thus obtain

$$\ll k^3 M^{3/2+\varepsilon_0/2+\varepsilon} X^{1/2} \ll k^3 M^{3/2+\varepsilon_0} M^{1/4} \ll k^3 M^{7/4+\varepsilon_0} \ll k^2 M^{2+\varepsilon_0},$$

since $k \ll M^{1/4}$.

In the case $k \gg \Delta M^{-1/2}$, we have $X \ll \Delta^4 M^{-1} k^{-6}$, and hence, the contribution is

$$\ll k^3 M^{3/2+\varepsilon_0/2+\varepsilon} X^{1/2} \ll k^3 M^{3/2+\varepsilon_0} (\Delta^4 M^{-1} k^{-6})^{1/2} \ll \Delta^2 M^{1+\varepsilon_0}.$$

6 Proof of Theorem 2

We begin by observing that we may assume M to be larger than a fixed large constant, because when $M \ll 1$, we also have $k \asymp \Delta \asymp V \asymp 1 \asymp M$ and the desired estimate for R reduces to $R \ll 1$, which would hold as certainly $R \ll 1 + M/V \ll 1$ in this case. Also, in the following all implicit constants are allowed to depend on δ and $\langle p, q \rangle$. We also make the simple observation that we may assume that $V \ll \sqrt{M}$ for the sums in question cannot obtain larger values by the Wilton–Jutila estimate.

Let $x \in [M, 2M]$, and let $N \in \mathbb{R}_+$ with $1 \ll N \ll M$. We will choose N later. The truncated Voronoi identity says that

$$\begin{aligned} & \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \\ &= \frac{k^{1/2}}{\pi \sqrt{2}} \sum_{n \leq N} a(n) n^{-3/4} e\left(-n \frac{\bar{h}}{k}\right) \\ & \quad \cdot \left((x + \Delta)^{1/4} \cos\left(4\pi \frac{\sqrt{n(x + \Delta)}}{k} - \frac{\pi}{4}\right) - x^{1/4} \cos\left(4\pi \frac{\sqrt{nx}}{k} - \frac{\pi}{4}\right) \right) \\ & \quad + O(k M^{1/2+\delta} N^{-1/2}). \end{aligned}$$

If x happens to be an integer, then the term $a(x) e(xh/k)$ is certainly $\ll x^\delta$ by Deligne’s estimate, and this is certainly $\ll k M^{1/2+\delta} N^{-1/2}$. Replacing the factor $(x + \Delta)^{1/4}$ by $x^{1/4}$ causes the error

$$\ll k^{1/2} \sum_{n \leq N} |a(n)| n^{-3/4} \Delta M^{-3/4} \ll k^{1/2} N^{1/4} \Delta M^{-3/4}.$$

Also, the difference of the cosines may be replaced by a sine integral:

$$\begin{aligned} & \cos\left(4\pi \frac{\sqrt{n(x + \Delta)}}{k} - \frac{\pi}{4}\right) - \cos\left(4\pi \frac{\sqrt{nx}}{k} - \frac{\pi}{4}\right) \\ &= - \int_x^{x+\Delta} \frac{2\pi \sqrt{n}}{k \sqrt{t}} \sin\left(4\pi \frac{\sqrt{nt}}{k} - \frac{\pi}{4}\right) dt. \end{aligned}$$

Combining the facts above gives

$$\sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right)$$

$$\begin{aligned}
&= -\sqrt{2} k^{-1/2} x^{1/4} \sum_{n \leq N} a(n) n^{-1/4} e\left(-n \frac{\bar{h}}{k}\right) \int_x^{x+\Delta} \sin\left(4\pi \frac{\sqrt{nt}}{k} - \frac{\pi}{4}\right) \frac{dt}{\sqrt{t}} \\
&\quad + O(k^{1/2} N^{1/4} \Delta M^{-3/4}) + O(k M^{1/2+\delta} N^{-1/2}).
\end{aligned}$$

We will split the interval $[M, 2M]$ into $\ll 1 + M/M_0$ closed subintervals of length at most $M_0 \in \mathbb{R}_+$ which we allow to have only endpoints in common. We shall choose the precise value of M_0 later. Also, we shall focus on one of the subintervals, say $J = [M, 2M] \cap [\alpha, \alpha + \Lambda]$, where $\alpha \in [M, 2M]$ and $\Lambda \in]0, M_0]$, which we assume to contain exactly $R_0 \in \mathbb{Z}_+$ of the original points x_1, \dots, x_R . Without loss of generality, we may assume these points to be x_1, \dots, x_{R_0} , ordered so that $x_1 < x_2 < \dots < x_{R_0}$. Once we have estimated R_0 from above as $\ll \Upsilon$, where Υ does not depend on J but only on k, M, Δ, δ and V , we can estimate R from above by

$$R \ll \Upsilon \left(1 + \frac{M}{M_0}\right).$$

Of course, if the subinterval contains none of the original points, then it trivially contains $\ll \Upsilon$ points.

Let us consider the choice of N in the truncated Voronoi identity. Provided that

$$N \ll M^3 V^4 \Delta^{-4} k^{-2} \quad \text{and} \quad N \gg k^2 M^{1+2\delta} V^{-2},$$

where the former implicit constant needs to be sufficiently small and the latter sufficiently large, the two error terms can be absorbed to the left-hand side, which in turn is $\gg V$, and we get for each $r \in \{1, \dots, R_0\}$ the estimate

$$\begin{aligned}
V &\ll \sum_{x_r \leq n \leq x_r + \Delta} a(n) e\left(n \frac{h}{k}\right) \\
&\ll k^{-1/2} M^{-1/4} \int_{x_r}^{x_r + \Delta} \left| \sum_{n \leq N} a(n) n^{-1/4} e\left(-n \frac{\bar{h}}{k}\right) \sin\left(4\pi \frac{\sqrt{nt}}{k} - \frac{\pi}{4}\right) \right| dt.
\end{aligned}$$

We shall actually choose N to be as small as possible, namely $N = c k^2 M^{1+2\delta} V^{-2}$ with a fixed constant $c \in \mathbb{R}_+$, though dependent on δ , and sufficiently large so that we can indeed absorb the term $k M^{1/2+\delta} N^{-1/2}$ to the left-hand side. We will have $N \ll M^{3-\delta} V^4 \Delta^{-4} k^{-2}$ since $V \gg k^{2/3} \Delta^{2/3} M^{-1/3+\delta}$, and so $N \ll M^3 V^4 \Delta^{-4} k^{-2}$ with a very small implicit constant, provided that M is sufficiently large. The requirement $N \ll M$ is satisfied thanks to the condition $V \gg k M^{2\delta}$. Similarly, the requirement

$N \gg 1$ is satisfied thanks to the condition $V \ll k M^{1/2+\delta}$. Also, we point out that, assuming that M is sufficiently large, we may assume that $N \geq 1$ for if $N < 1$, then $V \gg k M^{1/2+\delta}$, and we would have

$$k M^{1/2+\delta} \ll V \ll \sum_{x_1 \leq n \leq x_1 + \Delta} a(n) e\left(n \frac{h}{k}\right) \ll M^{1/2},$$

which is not possible for large M .

Next we cover the interval J with consecutive semiclosed intervals

$$I_1 = [\alpha, \alpha + V[, I_2 = [\alpha + V, \alpha + 2V[, \dots, I_\nu = [\alpha + (\nu - 1)V, \alpha + \nu V[,$$

where the number of intervals $\nu \in \mathbb{Z}_+$ is chosen so that it satisfies simultaneously the conditions $\nu \geq 2R_0$, $\nu > (M + \Delta)/V + 1$ as well as $\nu \ll M/V$. Let us temporarily simplify notation by writing

$$\Sigma(t) = \sum_{n \leq N} a(n) n^{-1/4} e\left(-n \frac{\bar{h}}{k}\right) \sin\left(4\pi \frac{\sqrt{nt}}{k} - \frac{\pi}{4}\right).$$

Let us consider integers

$$1 \leq a_1 < a_2 < a_3 < \dots < a_{R_0} \leq \nu$$

such that

$$x_1 \in I_{a_1}, \quad x_2 \in I_{a_2}, \quad \dots, \quad x_{R_0} \in I_{a_{R_0}},$$

and let $L = 1 + \lceil \Delta/V \rceil$ so that

$$[x_r, x_r + \Delta] \subseteq I_{a_r} \cup I_{a_{r+1}} \cup \dots \cup I_{a_{r+L}},$$

for each $r \in \{1, 2, \dots, R_0\}$. Furthermore, let $t_1 \in I_1, t_2 \in I_2, \dots, t_\nu \in I_\nu$ be points such that

$$|\Sigma(t_\ell)| = \max_{t \in I_\ell} |\Sigma(t)|$$

for each $\ell \in \{1, 2, \dots, \nu\}$. Now we may continue by estimating

$$V \ll k^{-1/2} M^{-1/4} \sum_{\ell=a_r}^{a_r+L} \int_{I_\ell} |\Sigma(t)| dt \ll k^{-1/2} M^{-1/4} \sum_{\ell=a_r}^{a_r+L} V |\Sigma(t_\ell)|.$$

Next, let us pick odd indices

$$1 \leq v_1 < v_2 < \dots < v_{R_0} \leq \nu$$

and even indices

$$2 \leq w_1 < w_2 < \dots < w_{R_0} \leq \nu$$

so that the absolute values $|\Sigma(t_{v_\ell})|$ for $\ell \in \{1, 2, \dots, R_0\}$ are the R_0 largest, counting multiplicities, among

$$|\Sigma(t_1)|, \quad |\Sigma(t_3)|, \quad |\Sigma(t_5)|, \quad \dots,$$

and similarly, so that the absolute values $|\Sigma(t_{w_\ell})|$ are the R_0 largest, counting multiplicities, among

$$|\Sigma(t_2)|, \quad |\Sigma(t_4)|, \quad |\Sigma(t_6)|, \quad \dots$$

Then we may continue our estimations by

$$\begin{aligned} R_0 &\ll k^{-1/2} M^{-1/4} \sum_{r=1}^{R_0} \sum_{\ell=a_r}^{a_r+L} |\Sigma(t_\ell)| \\ &\ll k^{-1/2} M^{\delta-1/4} \Delta V^{-1} \sum_{\ell=1}^{R_0} |\Sigma(t_{v_\ell})| + k^{-1/2} M^{\delta-1/4} \Delta V^{-1} \sum_{\ell=1}^{R_0} |\Sigma(t_{w_\ell})|, \end{aligned}$$

where the last estimate follows straightforwardly from the fact that the sums over ℓ intersect by at most $L+1$ terms and $L+1 \ll M^\delta \Delta V^{-1}$ thanks to the condition $V \ll \Delta M^\delta$. Without loss of generality and to simplify notation, we may assume that the term involving v_ℓ is larger, and we therefore can strike out here the terms involving w_ℓ , at the price of an extra constant factor 2. Now, by the Cauchy–Schwarz inequality,

$$R_0 \ll k^{-1/2} M^{\delta-1/4} \Delta V^{-1} \sqrt{R_0} \sqrt{\sum_{\ell=1}^{R_0} |\Sigma(t_{v_\ell})|^2},$$

so that

$$R_0 \ll k^{-1} M^{2\delta-1/2} \Delta^2 V^{-2} \sum_{\ell=1}^{R_0} |\Sigma(t_{v_\ell})|^2.$$

We split the sum $\Sigma(\cdot)$ dyadically, and write \sin in terms of $e(\pm \dots)$. Then

we continue by applying Bombieri's lemma, and estimating $\log M \ll M^\delta$,

$$\begin{aligned}
R_0 &\ll k^{-1} M^{2\delta-1/2} \frac{\Delta^2}{V^2} \sum_{r=1}^{R_0} \left| \sum_{n \leq N} \frac{a(n)}{n^{1/4}} e\left(-n \frac{\bar{h}}{k}\right) \sin\left(4\pi \frac{\sqrt{nt_{v_r}}}{k} - \frac{\pi}{4}\right) \right|^2 \\
&\ll k^{-1} M^{3\delta-1/2} \frac{\Delta^2}{V^2} \sum_{\pm} \sum_{r \leq R_0} \sum_{\substack{U \leq N/2 \\ \text{dyadic}}} \left| \sum_{U < n \leq 2U} \frac{a(n)}{n^{1/4}} e\left(-n \frac{\bar{h}}{k}\right) e\left(\pm \frac{2\sqrt{nt_{v_r}}}{k}\right) \right|^2 \\
&\ll k^{-1} M^{4\delta-1/2} \frac{\Delta^2}{V^2} \sum_{\pm} \max_{U \leq N/2} \sum_{r \leq R_0} \left| \sum_{U < n \leq 2U} \frac{a(n)}{n^{1/4}} e\left(-n \frac{\bar{h}}{k}\right) e\left(\pm \frac{2\sqrt{nt_{v_r}}}{k}\right) \right|^2 \\
&\ll k^{-1} M^{4\delta-1/2} \frac{\Delta^2}{V^2} \max_{U \leq N/2} U^{1/2} \max_{r \leq R_0} \sum_{s=1}^{R_0} \left| \sum_{U < n \leq 2U} e\left(\frac{2\sqrt{n}(\sqrt{t_{v_r}} - \sqrt{t_{v_s}})}{k}\right) \right|.
\end{aligned}$$

The terms with $s = r$ are easily seen to contribute

$$\ll k^{-1} M^{4\delta-1/2} \Delta^2 V^{-2} N^{3/2} \ll k^2 M^{1+7\delta} \Delta^2 V^{-5}.$$

To estimate the remaining terms, those with $s \neq r$, we first observe that

$$|\sqrt{t_{v_r}} - \sqrt{t_{v_s}}| \asymp \int_{t_{v_s}}^{t_{v_r}} \frac{dt}{\sqrt{t}} \asymp \frac{|t_{v_r} - t_{v_s}|}{M^{1/2}} \ll \frac{M_0}{M^{1/2}},$$

and so we may use the theory of exponent pairs to estimate

$$\begin{aligned}
&\sum_{U < n \leq 2U} e\left(\frac{2\sqrt{n}(\sqrt{t_{v_r}} - \sqrt{t_{v_s}})}{k}\right) \\
&\ll k^{-p} |\sqrt{t_{v_r}} - \sqrt{t_{v_s}}|^p U^{q-p/2} + \frac{k U^{1/2}}{|\sqrt{t_{v_r}} - \sqrt{t_{v_s}}|} \\
&\ll k^{-p} M_0^p M^{-p/2} U^{q-p/2} + \frac{k U^{1/2} M^{1/2}}{|t_{v_r} - t_{v_s}|}.
\end{aligned}$$

Thus, the remaining terms contribute, estimating again $\log M \ll M^\delta$ and remembering that $q \geq 1/2 \geq p$ so that $1/2 + q - p/2 > 0$,

$$\begin{aligned}
&\ll k^{-1} M^{4\delta-1/2} \Delta^2 V^{-2} \\
&\quad \cdot \max_{U \leq N/2} U^{1/2} \left(R_0 k^{-p} M_0^p M^{-p/2} U^{q-p/2} + \max_{r \leq R_0} \sum_{s \neq r} \frac{k U^{1/2} M^{1/2}}{|t_{v_r} - t_{v_s}|} \right) \\
&\ll k^{-1} M^{4\delta-1/2} \Delta^2 V^{-2} N^{q+1/2-p/2} R_0 k^{-p} M_0^p M^{-p/2} \\
&\quad + k^{-1} M^{5\delta-1/2} \Delta^2 V^{-2} N k M^{1/2} V^{-1} \\
&\ll R_0 \cdot k^{2q-2p} \Delta^2 M_0^p M^{q-p+\delta(5+2q-p)} V^{p-2q-3} + k^2 M^{1+7\delta} \Delta^2 V^{-5}.
\end{aligned}$$

We shall choose M_0 to be as large as possible so that the first term on the right-hand side will be $\ll R_0$ with a small implicit constant and can therefore be absorbed to the left-hand side. That is, we shall choose

$$M_0 \asymp k^{2-2q/p} \Delta^{-2/p} M^{1-q/p+\delta(1-2q/p-5/p)} V^{2q/p-1+3/p}.$$

Thus, we have estimated R_0 as

$$\ll k^2 M^{1+7\delta} \Delta^2 V^{-5}.$$

The total estimate for R is therefore

$$\begin{aligned} R &\ll k^2 M^{1+7\delta} \Delta^2 V^{-5} \left(1 + \frac{M}{M_0}\right) \\ &\ll k^2 M^{1+7\delta} \Delta^2 V^{-5} \\ &\quad + k^2 M^{1+7\delta} \Delta^2 V^{-5} M k^{2q/p-2} \Delta^{2/p} M^{-1+q/p+\delta(5/p+2q/p-1)} V^{-2q/p+1-3/p} \\ &\ll k^2 M^{1+7\delta} \Delta^2 V^{-5} + k^{2q/p} \Delta^{2+2/p} M^{1+q/p+\delta(6+5/p+2q/p)} V^{-2q/p-4-3/p}. \end{aligned}$$

7 Proof of Theorem 3

To estimate the integral

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^A dx,$$

we estimate it separately in the regions where the integrand is $\leq (M^{2\delta} V_0)^A$ and $\geq (M^{2\delta} V_0)^A$, where $\delta \in \mathbb{R}_+$ is small and fixed. The former values contribute $\ll M^{1+2\delta A} V_0^A$. To estimate the contribution from the latter values, we split the remaining value range dyadically into intervals of the shape $[V, 2V]$ with $V \in [V_0 M^{2\delta}, \infty[$. If necessary, we extend the last interval, losing at most a constant factor in the estimations. The number of subintervals is $\ll \log M \ll M^\varepsilon$. For each value interval, we choose a maximal number of points $x_1, \dots, x_{R(V)}$ from the interval $[M, 2M]$ so that

$$\left| \sum_{x_r \leq n \leq x_r+\Delta} a(n) e\left(n \frac{h}{k}\right) \right| \in [V, 2V]$$

for each $r \in \{1, \dots, R(V)\}$ and that $|x_r - x_s| \geq V$ for all $r, s \in \{1, \dots, R(V)\}$ with $r \neq s$. We recall that we certainly have $R(V) = 0$ if $V \gg \sqrt{M}$ or

$V \gg \Delta M^\delta$ or $V \gg k^\alpha \Delta^\beta M^{\gamma+2\delta}$. Now the contribution from the large values of the integrand is bounded by

$$\ll_A \sum_V V \cdot R(V) V^A,$$

where the summation over V is dyadic. Using Theorem 2, this is

$$\begin{aligned} \ll_\delta \sum_V & (k^2 M^{1+7\delta} \Delta^2 V^{-5} \\ & + k^{2q/p} \Delta^{2+2/p} M^{1+q/p+\delta(6+5/p+2q/p)} V^{-2q/p-4-3/p}) V^{A+1}. \end{aligned}$$

In each term V is estimated from below by V_0 or from above by $k^\alpha \Delta^\beta M^{\gamma+2\delta}$, depending on whether the final exponent of V is negative or positive. Upon letting δ have smaller and smaller values, the first term in the parentheses gives rise to Φ and the second to Ψ .

8 Proof of Theorem 4

Proof of Theorem 4. We choose $p = q = 1/2$. By Theorem 5.5 in [6]

$$\sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \ll \Delta^{1/6} M^{1/3+\varepsilon} \ll M^{5/(12.6)} M^{1/3+\varepsilon} \ll M^{29/72+\varepsilon}.$$

Notice that when $k \gg M^{1/9}$, this bound is superior to $k^{1/4} M^{3/8+\varepsilon}$ from [24]. Hence using Theorem 3, we obtain, for any $V_0 \in [1, \infty[$ with $k \ll V_0 \ll M^{29/72}$ and $V_0 \gg k^{2/3} \Delta^{2/3} M^{-1/3}$, that

$$\begin{aligned} \int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) e\left(n \frac{h}{k}\right) \right|^A dx \\ \ll M^{1+\varepsilon} V_0^A + k^2 M^{11/6+29(A-4)/72+\varepsilon} + k^2 M^{9/2+29(A-11)/72+\varepsilon}. \end{aligned}$$

The term V_0 does not appear anywhere else except in the main term, so we can choose it to be as small as possible, namely k . For this choice, we also have $V_0 \gg k^{2/3} M^{-1/18} = k^{2/3} \Delta^{2/3} M^{-1/3}$. The contribution of the main term is $k^A M^{1+\varepsilon}$. The three terms satisfy

$$k^2 M^{11/6+29(A-4)/72+\varepsilon} \gg k^2 M^{9/2+29(A-11)/72+\varepsilon},$$

and

$$k^2 M^{11/6+29(A-4)/72+\varepsilon} \gg k^A M^{1+\varepsilon},$$

and we get the claimed bound. \square

9 Proof of Theorem 5

Proof of Theorem 5. We will first apply Theorem 14 with the exponent pair $p = 4/18$ and $q = 11/18$ and the parameters $\alpha = \gamma = \delta = 0$, $k = 1$ and $\beta = 1/3$. Now the main term becomes $M^{A/4+1}$. The term Φ becomes $M^{(A+1)/3+\varepsilon}$ and the term Ψ becomes $M^{A/4+1+\varepsilon}$. Since $(A+1)/3 < A/4+1$ exactly when $A < 8$, we have now derived

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) \right|^A dx \ll \begin{cases} M^{A/4+1+\varepsilon} & \text{when } A \leq 8, \\ M^{(A+1)/3+\varepsilon} & \text{when } A \geq 8. \end{cases}$$

We will now apply Theorem 3 with exponent pair $p = q = 1/2$. By the trivial estimate and by the estimate for a long sum, we know that

$$\sum_{x \leq n \leq x+\Delta} a(n) \ll \min(\Delta M^\varepsilon, M^{1/3+\varepsilon}).$$

Using these bounds we obtain

$$\begin{aligned} & \int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) \right|^A dx \\ & \ll \begin{cases} M^{1+\varepsilon} V_0^A + \Delta^{A-2} M^{1+\varepsilon} + \Delta^6 M^{2+\varepsilon} V_0^{A-11} & \text{when } \Delta \ll M^{1/3}, \\ M^{1+\varepsilon} V_0^A + \Delta^2 M^{(A-1)/3+\varepsilon} + \Delta^6 M^{2+\varepsilon} V_0^{A-11} & \text{when } \Delta \gg M^{1/3}. \end{cases} \end{aligned}$$

Let us now choose V_0 so that the first and the last term are the same (up to an epsilon):

$$M V_0^A = \Delta^6 M^2 V_0^{A-11},$$

which is equivalent with $V_0 = \Delta^{6/11} M^{1/11}$. Clearly, this choice satisfies $V_0 \gg 1 = k$ and is easily seen to satisfy $V_0 \gg \Delta^{2/3} M^{-1/3}$. It also satisfies $V_0 \ll \Delta M^\varepsilon$ and $V_0 \ll M^{1/3+\varepsilon}$ since $M^{1/5} \ll \Delta \ll M^{4/9}$. The estimate becomes now

$$\begin{aligned} & \int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) \right|^A dx \\ & \ll \begin{cases} M^{1+A/11+\varepsilon} \Delta^{6A/11} + \Delta^{A-2} M^{1+\varepsilon} & \text{when } \Delta \ll M^{1/3}, \\ \Delta^2 M^{(A-1)/3+\varepsilon} + M^{1+A/11+\varepsilon} \Delta^{6A/11} & \text{when } \Delta \gg M^{1/3}. \end{cases} \end{aligned}$$

When $\Delta \ll M^{1/3}$, we have $\Delta^{5A/11-2} \ll M^{A/11}$ and hence $M^{A/11} \Delta^{6A/11} \gg \Delta^{A-2}$. Thus, $M^{1+A/11+\varepsilon} \Delta^{6A/11} \gg \Delta^{A-2} M^{1+\varepsilon}$.

When $\Delta \gg M^{1/3}$, we have $\Delta \gg M^{1/3} \gg M^{(4A-22)/(9A-33)}$, so that $\Delta^2 M^{(A-1)/3+\varepsilon} \ll \Delta^{6A/11} M^{1+A/11+\varepsilon}$. We have now derived

$$\int_M^{2M} \left| \sum_{x \leq n \leq x+\Delta} a(n) \right|^A dx \ll M^{1+A/11+\varepsilon} \Delta^{6A/11}.$$

Finally, the proof is completed by comparing the above bounds separately in the cases $A \geq 8$ and $A \leq 8$. \square

Acknowledgements

The first author was funded by the Academy of Finland project 138337, by the Finnish Cultural Foundation, and by the Ruth och Nils-Erik Stenbäcks stiftelse. The second author was funded by the Academy of Finland through the Finnish Centre of Excellence in Inverse Problems Research and the projects 276031, 282938, 283262 and 303820, by the Magnus Ehrnrooth Foundation, by the Finnish Cultural Foundation, by the Foundation of Vilho, Yrjö and Kalle Väisälä, and by the Basque Government through the BERC 2014–2017 program and by Spanish Ministry of Economy and Competitiveness MINECO: BCAM Severo Ochoa excellence accreditation SEV-2013-0323.

References

- [1] CRAMÉR, H.: *Über zwei Sätze von Herrn G. H. Hardy*, Math. Z., 15 (1922), 200–210.
- [2] DELIGNE, P.: *La conjecture de Weil: I*, Publ. Math. Inst. Hautes Études Sci., 43 (1974), 273–307.
- [3] ERNVALL-HYTÖNEN, A.-M.: *On the error term in the approximate functional equation for exponential sums related to cusp forms*, Int. J. Number Theory, 4 (2008), 747–756.
- [4] ERNVALL-HYTÖNEN, A.-M.: *On the mean square of short exponential sums related to cusp forms*, Funct. Approx. Comment. Math., 45 (2011), 97–104.
- [5] ERNVALL-HYTÖNEN, A.-M.: *Mean square estimate for relatively short exponential sums involving Fourier coefficients of cusp forms*, Ann. Acad. Sci. Fenn. Math., 40 (2015), 385–395.
- [6] ERNVALL-HYTÖNEN, A.-M., and K. KARPPINEN: *On short exponential sums involving Fourier coefficients of holomorphic cusp forms*, Int. Math. Res. Not., article ID rnn022 (2008), 1–44.
- [7] GRAHAM, S. W., and G. KOLESNIK: *Van der Corput’s Method of Exponential Sums*, London Mathematical Society Lecture Note Series, 126, Cambridge University Press, 1991.

- [8] IVIĆ, A.: *Large values of the error term in the divisor problem*, Invent. Math., 71 (1983), 513–520.
- [9] IVIĆ, A.: *On the divisor problem and the Riemann zeta-function in short intervals*, Ramanujan J., 19 (2009), 207–224.
- [10] IVIĆ, A., and P. SARGOS: *On the higher moments of the error term in the divisor problem*, Illinois J. Math., 81 (2007), 353–377.
- [11] IVIĆ, A., and W. ZHAI: *On the Dirichlet divisor problem in short intervals*, Ramanujan J., 33 (2014), 447–465.
- [12] JÄÄSAARI, J., and E. V. VESALAINEN: *Exponential sums related to Maass forms*, Acta Arith., 190 (2019), 1–48.
- [13] JUTILA, M.: *On the divisor problem for short intervals*, Ann. Univ. Turku. Ser. A I, 186 (1984), 23–30.
- [14] JUTILA, M.: *On exponential sums involving the divisor function*, J. Reine Angew. Math., 355 (1985), 173–190.
- [15] JUTILA, M.: *Lectures on a Method in the Theory of Exponential Sums*, Lectures on Mathematics, 80, Tata Institute of Fundamental Research, 1987.
- [16] JUTILA, M.: *On exponential sums involving the Ramanujan function*, Proc. Indian Acad. Sci., 97 (1987), 157–166.
- [17] JUTILA, M., and Y. MOTOHASHI: *Uniform bound for Hecke L-functions*, Acta Math., 195 (2005), 61–115.
- [18] MONTGOMERY, H. L.: *Topics in Multiplicative Number Theory*, Lecture Notes in Mathematics, 227, Springer-Verlag, 1971.
- [19] RANKIN, R. A.: *Contributions to the theory of Ramanujan’s function $\tau(n)$ and similar arithmetical functions II. The order of Fourier coefficients of integral modular forms*, Math. Proc. Cambridge Philos. Soc., 35 (1939), 357–372.
- [20] ROBERT, O., and P. SARGOS: *Three-dimensional exponential sums with monomials*, J. Reine Angew. Math., 591 (2006), 1–20.
- [21] SELBERG, A.: *Bemerkungen über eine Dirichletsche Reihe, die mit der Theorie der Modulformen nahe verbunden ist*, Arch. Math. Naturvid., 43 (1940), 47–50.
- [22] TANIGAWA, Y., and W. ZHAI: *On the fourth power moment of $\Delta(x)$ and $E(x)$ in short intervals*, Int. J. Number Theory, 5 (2009), 355–382.
- [23] TSANG, K.-M.: *Higher-power moments of $\Delta(x)$, $E(t)$ and $P(x)$* , Proc. London Math. Soc., 65 (1992), 65–84.
- [24] VESALAINEN, E. V.: *Moments and oscillations of exponential sums related to cusp forms*, Math. Proc. Cambridge Philos. Soc., 162 (2017), 479–506.
- [25] WILTON, J. R.: *A note on Ramanujan’s function $\tau(n)$* , Math. Proc. Cambridge Philos. Soc., 25 (1929), 121–129.
- [26] WU, J., and W. ZHAI: *Distribution of Hecke eigenvalues of newforms in short intervals*, Q. J. Math., 64 (2013), 619–644.